AWARD NUMBER: W81XWH-14-1-0500

TITLE: Toward a Molecular Understanding of Noise-Induced Hearing Loss

PRINCIPAL INVESTIGATOR: Ronna Hertzano, MD PhD

CONTRACTING ORGANIZATION: University of Maryland, Baltimore

Baltimore, MD 21201

REPORT DATE: October 2017

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command

Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;

Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

## REPORT DOCUMENTATION PAGE

Form Approved OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

October 2017	Annual	30Sep2016-29Sep2017
4. TITLE AND SUBTITLE	Annual	5a. CONTRACT NUMBER
4. IIILE AND SUBTILE		5a. CONTRACT NUMBER
		EL ODANTAUMDED
Toward a Molecular Understa	anding of Noise-Induced Hearing Loss	5b. GRANT NUMBER
		W81XWH-14-1-0500
		5c. PROGRAM ELEMENT NUMBER
6. AUTHOR(S)		5d. PROJECT NUMBER
		F. TAOK NUMBER
Ronna Hertzano, MD, PhD		5e. TASK NUMBER
		5f. WORK UNIT NUMBER
F Mail alors and Romail and Inc.	1 . 1	SI. WORK UNIT NUMBER
E-Mail: rhertzano@smail.umaryland		A DEDECRMING ORGANIZATION DEPORT
7. PERFORMING ORGANIZATION NAME(S		8. PERFORMING ORGANIZATION REPORT NUMBER
University of Maryland, Baltimore, MD 21201	ltimore	NOMBER
Baltimore, MD 21201		
9. SPONSORING / MONITORING AGENCY	NAME(S) AND ADDRESS(ES)	10. SPONSOR/MONITOR'S ACRONYM(S)
U.S. Army Medical Research and M	lateriel Command	
Fort Detrick, Maryland 21702-5012		11. SPONSOR/MONITOR'S REPORT
· •		NUMBER(S)
40 DIOTDIDUTION / AVAIL ADULTY OTATE		

#### 12. DISTRIBUTION / AVAILABILITY STATEMENT

Approved for Public Release; Distribution Unlimited

#### 13. SUPPLEMENTARY NOTES

#### 14. ABSTRACT

The purpose of this project is to generate the cell type-specific molecular blueprint of changes in gene expression following different types of noise exposure and their treatments, in the inner ear. To this end, we have (a) Established the hair cell (HC) and supporting (SC) cell-specific translatome of adult mouse inner ears; (b) Established the molecular changes induced by PTS-resulting noise exposure in HC, SC and whole inner ears, 6 and 24 hours after noise exposure and began the TTS analysis; (c) We identified critical differences in the response of male and female mice to noise, and obtained approval to modify Specific Aim 3 to focus on the response to Heat Shock profiling male and female mice, separately.

#### 15. SUBJECT TERMS

Permanent threshold shift, Temporary threshold shift, Noise induced hearing loss, Ribotag, RNA-seq, hair cell, supporting cell, SAHA, Heat shock, sex differences

16. SECURITY CLASS	SIFICATION OF:		17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT	b. ABSTRACT	c. THIS PAGE	Unalogoified	12	19b. TELEPHONE NUMBER (include area code)
Unclassified	Unclassified	Unclassified	Unclassified	12	,

Standard Form 298 (Rev. 8-98) Prescribed by ANSI Std. Z39.18

		<u>Page</u>
	Table of Contents	
1.	Introduction	4
2.	Keywords	4
3.	Accomplishments	4
4.	Impact	9
5.	Changes/problems	10
6.	Products	10
7.	Participants and other collaborating organizations	10
8.	Special reporting requirements	12
9.	Appendices	N/A

#### 1. INTRODUCTION

Noise induced hearing loss (NIHL) is a major health concern for the Department of Defense. Noise exposure often is inevitable, and may results in a permanent loss of hearing. Unfortunately, there are no treatments to prevent or reverse NIHL. As a first step towards designing targeted therapeutics, we suggested to generate mouse models which allow cell type-specific translatome analysis in the ear. These, in turn, will be used to analyze the genes expressed in the hair cells (HC) and supporting cells (SC) of adult mice before and after different types of noise exposure as well as pre-conditioning treatments, which in mice, can ameliorate NIHL. Here we report our progress over the third year of the project, in which (a) HC and SC from adult mice were sequenced and their expression patterns were compared and analyzed; (b) we established the molecular cell type-specific blueprint following PTS-inducing noise exposure; (c) we identified differences in the molecular response to PTS- and TTS-inducing noise exposure; and (d) we identified critical differences in the response of male and female mice to noise exposure, which lead to modification of Specific Aim 3 (manuscript in preparation, data shared at the ARO in an oral presentation).

#### 2. KEYWORDS

Permanent threshold shift, Temporary threshold shift, Noise induced hearing loss, Ribotag, RNA-seq, Hair cell, Supporting cell, SAHA, Heat shock, Sex differences.

#### 3. ACCOMPLISHMENTS


<u>Specific Aim 1</u>: To determine the OHC- and SC-specific transcriptional and signaling cascades activated in vivo in response to PTS-inducing noise exposure

- Major Task 1: To establish the OHC- and SC-specific translatome of adult mouse inner ears. Progress by subtasks:
  - i. Obtain ACURO approval following UMSOM IACUC approval complete.
  - ii. Mouse crosses and tissue harvesting complete.
  - iii. Tissue processing complete, polysome IP complete, submission of samples for RiboTag-seq complete; RiboTag-seq complete.
- We identified 410 genes with a significantly higher expression in OHC compared with SC and the rest of the inner ear. Many of the genes have not been previously described and/or studied in relation to inner ear development/function. Within the top 10 genes we find: Klhdc7b, Strc, Dlec1, Btn2a2, Slc26a5, Gfi1, Lmod3, Ocm and Lhfpl5. Of these

ten genes, six (marked in bold) are known to be expressed in hair cells and when mutated to cause a deafness phenotype. Many additional deafness genes are included in the list, indicating that their neighboring genes are highly likely to have a critical role in maintenance of the hair cells. Importantly, Scl26a5, which was used to pull down the ribosomes in this assay is in the top 5 most enriched genes, validating our approach. A cis-regulatory analysis identified IKZF2 and GFI1 as the likely most prominent regulators of the adult OHC-translatome. While IKZF2 role in hearing has yet to be published, in a separate project we identified IKZF2 as a critical regulator of OHC electromotility, further validating our findings. Two manuscripts are now in preparation.

- We also identified 282 SC enriched genes. Much less is known about the adult supporting cell translatome. The following genes were within the top 10 most enriched SC genes: Igfbp6, Bpifa1, Timp4, Aldoc, Ascbg1, Gldc, Egfl6, Plbd1, Mlc1 and Slc25a18. From these, Bpifa1 has been previously associated with susceptibility to otitis media (PMID 25765466), Timp4 with worse outcomes after myocardial infarction in mice (PMID 20516072), Aldoc was previously described as expressed in the ear but at a very low resolution and only in neonatal mice (PMID 24475166), and Slc25a18 has a role with Glutamate transport. Importantly, Sox2, which was used to pull down the ribosomes, is within the top 30 enriched genes. Its lower location compared to Slc26a5 highlights how little we know about SC in the adult mouse. Also, many of the enriched genes, do follow even in the early postnatal mouse a pattern of increasing expression in the non-HC population based on the SHIELD further increasing our confidence level with regards to their specific expression in adult tissues. Finally, a cis-regulatory analysis of the SC-enriched genes pointed to SOX2 and REST as their likely key regulators.
- <u>Specificity of cre-drivers</u>. An important step in the data analysis of each experiment is careful characterization of its individual components. Here we validated the specificity of the cre-drivers for OHC and SC (see **Figure 1** and **Table 1**). The OHC-cre, driven by the prestin locus was found to be specific for OHC (the red reported is shown only in OHC,

Figure 1) and HC and OHC transcripts are enriched in the immunoprecipitate while all other transcripts are depleted (Table 1). The SC-cre, driven by the Sox2 gene was found to induce recombination both in SC and in glia (see Figure 1 and Table 1). This result was not anticipated, however, it does not take away from the validity of the data. We now know that SC transcripts in this dataset include also glia.



Sox2CreERT2 reporter – expression in supporting cells as well as glia throughout the cochlear duct.

**Figure 1 Validation of OHC and SC Cre drivers.** PrestinCreERT2 (credriver for OHC, left panel) and Sox2CreERT2 (driver for SC, right panel) were crossed with a mouse that expresses a red fluorescent protein in the presence of cre-recombinase (Ai14).

		Rpl2	2 <sup>HA/+</sup> ;Prest	in <sup>CreERT2/+</sup>	Rpl	22 <sup>нд/+</sup> ;Sох	2 <sup>CreERT2</sup> /+
Cell Types	Gene Symbol	Input	IP	Enrichment /Depletion (Log2)	Input	IP	Enrichment /Depletion (Log2)
Outer Hair	Slc26a5	134	559	2.06	414	144	-1.53
Cells	Ocm	793	5624	2.83	2061	1316	-0.65
Hair Cells	Gfi1	53	200	1.91	136	44	-1.63
Hair Cells	Strc	128	763	2.57	388	141	-1.47
Supporting	Sox2	336	186	-0.85	544	2984	2.46
Cells	Jag1	444	431	-0.04	847	2729	1.69
Nourons	Tubb3	1453	608	-1.26	3407	3820	0.16
Neurons	Dlg4	1165	979	-0.25	3984	5779	0.54
Glia	Gfap	165	101	-0.71	1212	6639	2.45
(Astrocytes)	Aldh1l1	327	209	-0.64	1198	2422	1.02
Masonshima	Pou3f4	2023	993	-1.03	5501	1523	-1.85
Mesenchyme	Etv1	546	242	-1.18	1464	728	-1.01

Table 1 - Enrichment analysis of immunoprecipitated transcripts from the Prestin and Sox2 cre mice.

- Major Task 2: To determine the OHC- and SC-specific transcriptional and signaling cascades activated in response to PTS-inducing noise injury. Progress by subtasks:
  - i. Mouse crosses, noise exposure, tissue harvesting, histological analysis, ABR and DPOAE measurements. Complete.
  - ii. Tissue processing complete.
  - iii. Data analysis largely reported last year.
  - iv. Validation experiments polysome IP to be used for RT-qPCR complete. Tissue harvesting for immunohistochemistry and in situ hybridization complete. Validation experiments to be performed with NanoString in the last quarter of year IV of the project.

#### Future plans:

- Validation of select genes by in-situ hybridization and/or immunohistochemistry
- Possible interrogation of select mouse mutants for these genes (as part of this or future applications)


# <u>Specific Aim 2</u>: To determine the OHC- and SC-specific signaling cascades activated in vivo in response to otoprotective interventions.

This Aim was designed to define the cell type-specific molecular blueprint of interventions that may ameliorate NIHL. We planned to characterize TTS-inducing noise exposure, heat shock and restraint stress. In our previous annual reports we discussed the rationale for maintaining the TTS and Heat Shock. The heat shock goal was transferred to a modified Specific Aim 3 that is also sex-specific.

<u>TTS-inducing noise exposure</u>: crosses, calibration, validation cytocochleograms, noise exposure, tissue harvesting, polysome IP and RNA extraction was completed both for the OHC and SC crosses. Sequencing (of tissue obtained from males and females, combined) is complete. Analysis – ongoing. <u>Preliminary Results</u>

In the analysis of tissue obtained from whole ears (IP tissue has been sequenced but analysis is ongoing) we identified 12 clusters of genes that are distinct in their behaviors (Figure 2). For

example, cluster 1 consists of 176 genes that are uniquely downregulated following TTSinducing noise exposure. In contrast cluster 3 consists of 49 genes that are uniquely upregulated after TTSinducing noise exposure. As TTSinducing noise exposure has also been shown to be otoprotective from PTS, chemotherapy and aminoglycoside exposure - these genes are well positioned to be candidate otoportective genes. Interestingly, within this group we find DFNA5, which in its absence mice and human suffer from sensorineural hearing



**Figure 2 cluster analysis of the differences between the molecular response to PTS and TTS**. In each graph the first point represents the baseline values, followed by expression at 6 and 24 hours, first in PTS and then in TTS. Y axis - log2 transformed.

loss, further supporting our hypothesis. **Cluster 2** consists of genes that increase within 24 hours of exposure to PTS inducing noise exposure but not after TTS inducing noise exposure. This cluster is statistically significantly enriched for genes that encode proteins of the innate immune response (p-value 3.65e-06 and corrected p-value 0.009) (see **Figure 3**). The significance of this finding is that one of the defining features differentiating between the response to PTS and TTS is activation of the innate immune response, a finding with a clear translational significance.

An interesting observation as part of the results of both specific aims 1 and 2 is an activation of 'immediate early genes' also known as 'stress genes' in the samples exposed to noise (e.g., Fos, Jun, Hspa1a). The increase in gene expression seen at 6 hours decreases dramatically for some of the genes at 24 hours and for other genes persists (see **Figure** 

innate immune response - GO:0045087 Raw p-value: 3.65e-06

Corrected p-value: 0.00900


Number of genes: 12.0 Frequency in set: 16.09

Gene ID	Gene symbol	Probe ID
12260	Ciqb	(ENSMUSGooocoo)
232371	Cırl	(ENSMUSG0000003.
230073	Ddx58	(ENSMUSG0000004.
14469	Gbp2	(ENSMUSGooocoo)
71586	lfih1	(ENSMUSG0000002.
15957	lfit1	(ENSMUSG0000003.
66141	lfitm3	(ENSMUSG0000002.
54123	Irf7	(ENSMUSG0000002
15944	lrgm1	(ENSMUSG0000004
20684	Sp100	(ENSMUSG0000002
20846	Stat1	(ENSMUSG0000002
142980	TIra	rENSMUSG0000003.

Figure 3 Innate immune response genes are enriched in cluster 2.

**4**). A recent publication (Nature Methods, Vol 14 No.10, October 2017) showed that this particular group of genes is activated in a non-specific manner following tissue dissociation (**Figure 4**). This

finding further strengthens the importance of our chosen approach for interrogating gene expression following noise exposure – as cell sorting or even just tissue dissociation for traditional single-cell RNA-seq is likely to invoke numerous changes in gene expression that would specifically mask changes of interest in this type of experiments.



Specific Aim 3: Following the in-person review in the DoD this summer, specific aim 3 was redesigned to allow us to both study

Figure 4 Overlap between dissociation and noise-induced changes in gene expression

the response to Heat Shock with and without noise exposure as originally planned as well as obtain all tissue separately from male and female mice. This was presented, submitted to and approved by the DoD. We are currently in the process of tissue collection and hope to complete that stage by March 2018.

## **Key research accomplishments**

- Identification of OHC and SC unique genes and their regulators.
- Molecular description of the response to PTS in OHC, SC and whole inner ear.
- Molecular description of the response to TTS.
- Key differences between the response to TTS and PTS.
- Critical differences in the response to noise between males and females.

## **Conclusion**

The results of the cell type-specific molecular changes as a result of PTS-inducing noise exposure will set a foundation for rationale design of drugs to treat NIHL. This is critically important for the military, hearing loss and tinnitus are a principal cause for disability in the DoD and to date there are no efficient preventative or curative treatments. Maximal benefit from our molecular data is pending the TTSinducing noise exposure dataset, as the latter will allow to distinguish protective changes in gene expression from those that represent a detrimental effect on hearing. Indeed, our initial analysis shows that at least one deafness causing gene (DFNA5) is upregulated specifically after TTS-resulting noise exposure. In addition, our results demonstrate the feasibility of using cell type-specific molecular changes as a readout in future studies to protect from NIHL, and the advantage of the RiboTag approach over the single-cell or dissociation-based approaches. With our current results we anticipate being able to develop a molecular panel predictive of type of noise injury that could be used as a fast-readout to screen for the effect of therapeutics. Our identified differences in the physiologic response to noise between male and female mice are important, not only for the experimental design of future studies, but as both men and women now serves in the army in positions where they suffer from NIHL, and we should therefore design treatments that will be efficacious in both sexes. In addition, the new structure of specific aim 3 will enable us to understand, at a molecular level, the differences between the response of male and female mice to noise.

#### Publications, abstracts, and presentations

- Three manuscripts are in the final stages of preparation for publication: the kit comparison manuscript, differences in the response to noise between male and female mice, and the identification of IKZF2 as a key regulator of the OHC translatome.
- The findings about the sex-differences, which are critically important for experimental design were shared at the ARO 2017 Mid-Winter Meeting (pre-publication, oral presentation)
  - Mitra S, Drake V, Margulies Z, Milon B, Song Y, Depireux D and Hertzano R (2017) The impact of sex on the response to noise and otoprotective therapies against acoustic injury in mice; Association for Research in Otolaryngology, Baltimore, MD, USA.
- The findings of the kit-comparison work were shared at the ARO (pre-publication, oral presentation)
  - Song Y, Milon B, Ott S, Zhao X, Sadzewicz L, Mahurkar A and Hertzano R (2017) RiboTag-Seq: a comparative analysis of library prep approaches for sequencing low

input translatome samples; Association for Research in Otolaryngology, Baltimore, MD, USA.

## Opportunities for training and professional development

<u>Training or fellowship awards</u>:

Sunayana Mitra, PhD, participated in the Association for Research in Otolaryngology meeting, as well as in the EARssentials course at the NIH.

Beatrice Milon, PhD, participated in the Association for Research in Otolaryngology meeting.

Ryan Casserly, MD, worked in the Hertzano laboratory in a 5-month full time research rotation to characterize the differential response to noise between male and female mice.

Laboratory meetings - since obtaining funding from the DoD the entire Hertzano laboratory engages in in-depth study of current literature and techniques to study NIHL and has been increasing their knowledge and experience through laboratory meetings and journal clubs. The team has also trained two additional laboratories in the department (laboratories of Drs. Ahmed and Riazuddin) who now focus some of their work on NIHL. In addition, this past year, we developed a new series of laboratory meetings named H&H (Hearing and Hormones) attended by Dr. Jessica A. Mong (a neuroendocrinologist with a focus on sex differences in the brain) and our group.

## Professional development

All members working on the DoD project participate in laboratory meetings, the translational Auditory and Vestibular research day, the Association for Research in Otolaryngology mid-winter meeting.

## Inventions, patents and licences

Nothing to report

#### Reportable outcomes

Nothing to report

## Other achievements

Nothing to report.

## References

## 4. IMPACT

- Our work and its presentation in scientific meetings has significantly increased awareness in the ear field to cell type specific molecular analysis using the RiboTag mice, and better approaches for sequencing of their translatomes.
- Our work and its presentation in scientific meetings has brought sex differences in the response to noise exposure to the awareness of the ear field scientific community and we believe will have broad impact on study designs.
- The results of our work will form the foundation for new rationale design of therapeutics to prevent noise induced hearing loss.

## 5. **CHANGES/PROBLEMS**

We had a delay in the breeding efficiency this year and therefore the baseline heat shock experiments are done on C57BL/6 and not the mixed background. However, with improvements to the sequencing protocol, we can now catch up over the year thanks to requiring less mice per biological replicate.

All changes in structure and expenditures have been reported in the in-person review (within the presentation) and are within the limits of the budget of the project.

No significant changes were made to biohazards.

## 6. PRODUCTS

Nothing to report.

#### 7. PARTICIPANTS AND OTHER COLLABORATING ORGANIZATIONS

## Individuals who work on the project

Name	Ronna Hertzano
Project Role	PI
Researcher identifier	
Nearest person month worked:	2
Contribution to the project	Overall responsibility for the proposal and all aspects of the research program including: hiring and training personnel, ensuring quality of data, interpretation of data, oversight of methods, administrative responsibility and reporting to the DoD.
Funding support	NIH R01, DC013817; Action on Hearing Loss. G65_Bowl; NIH R01, DC003544; Hearing Health Foundation – HRP support for gEAR.

Name	Didier Depireux
Project Role	Co-I
Researcher identifier	
Nearest person month worked:	1.2
Contribution to the project	Oversight of the noise exposure protocols, ABR and DPOAE setup and measurements; Discussion and analysis of Male/Female data.
Funding support	MII, Translational Research in Hearing Foundation, Capita foundation, NIH/NIDCR

Name	Ran Elkon
Project Role	Co-I
Researcher identifier	
Nearest person month	1.2
worked:	

Contribution to the	Data analysis and study design
project	
Funding support	

Name	Yang Song
Project Role	Analyst
Researcher identifier	
Nearest person month worked:	1.2
Contribution to the project	Data analysis
Funding support	

Name	Beatrice Milon
Project Role	Research Supervisor
Researcher identifier	
Nearest person month worked:	6 months
Contribution to the project	Study design, tissue collection, schedule oversight, training Yoko Ogawa, RiboTag IP, RNA analysis, cytocochleograms
Funding support	NIH R01, DC013817

Name	Yoko Ogawa
Project Role	Post Doctoral Fellow
Researcher identifier	
Nearest person month worked:	5 months
Contribution to the project	Tissue collection, cytocochleograms, validation (currently setting up), animal care
Funding support	

Name	Ryan Casserly
Project Role	Resident
Researcher identifier	
Nearest person month worked:	3 months
Contribution to the project	Characterized the sex differences in hearing
Funding support	

Name	Sunayana Mitra
Project Role	Post Doctoral Fellow
Researcher identifier	

Nearest person month worked:	12 months
Contribution to the	Performs all noise exposures, hearing measurements, pre-conditioning,
project	tissue collection, animal care, assists with writing animal protocols
Funding support	

## Changes in the other support of the PI/ other key personnel:

No significant changes.

#### 8. SPECIAL REPORTING

## **QUAD CHART**

